

## *Reflex Testing*

Reflex testing refers to laboratory tests which are automatically performed based on predetermined criteria. These tests will be performed unless Alomere Health Hospital is notified otherwise. The following is a list of Alomere's reflex testing:

| <b>Test Ordered</b>                         | <b>Reflex Test<br/>(If Ordered Test is Positive or<br/>Otherwise Indicated)</b>   |
|---|---|
| Antibody Screen                             | Antibody identification, antibody titer, and antigen type units if applicable.  |
| Type and Screen                             | If antibody screen is positive. 2 units of leuko-reduced packed RBCs are crossmatched   |
| Bacterial Culture                           | Bacterial identification and minimum inhibitory concentration (MIC) will be performed when indicated.   |
| Blood Culture                               | If test positive, Eplex Identification BCID-GP or BCID-GN   |
| CBC   | Peripheral smear and interpretation performed when indicated by pathologist   |
| Direct Antiglobulin Test Positive           | Elution   |
| Fetal Maternal Hemorrhage (FMH) Screen      | If positive, specimen is referred for #88841 "Fetomaternal Bleed, Flow Cytometry, Blood" for determination of RhIg dosage.                            |
| Fungal Culture                              | Fungal identification if indicated  |
| <i>Giardia/Cryptosporidium</i> Antigen      | Tests are performed and reported simultaneously when either is ordered.   |
| Hepatitis B Surface Antigen                 | Hepatitis B Surface Antigen Confirmation  |
| Hepatitis C Antibody                        | HCV RNA Detection Quant by PCR  |
| HIV Antibody                                | HIV Antibody Confirmation and Differentiation HVDIP   |
| Human Papillomavirus (HPV), high risk (PCR) | Human Papillomavirus Types 16 & 18/45 only (if appropriate)   |
| Influenza Antigen                           | Both Influenza A and B are performed  |
| Lactic Acid (Inpatient only)                | This test will generate a REFLEX Lactic Acid to be drawn Q2H after initial testing was collected. If the reflex is not needed, please notify the lab. |
| Lyme Disease Screen                         | <i>Borrelia Burgdorferi</i> AB, Immunoblot LYWB   |
| Pathology Examinations                      | ERA/PRA, FISH, Her2neu, or any additional assays as appropriate for findings.   |
| Treponemal Antibody IgG/IgM                 | <i>Treponema Pallidum</i> Ab, Total, with Reflex to confirmation SYPH1  |
| Troponin I (Inpatients only)                | This test will generate a REFLEX Troponin I to be drawn Q2H after initial testing was collected. If the reflex is not needed, please notify the lab.  |